arXiv:nucl-th/9704046v1 22 Apr 1997

# Strangeness : Theoretical Status*

J. Cleymans
Department of Physics, University of Cape Town, South Africa

February 9, 2008

**Abstract**

Strangeness plays an important role in the study of quark matter since it indicates the rate at which new particles are being produced and therefore provides information about the degree of chemical equilibration reached in heavy ion collisions. It is shown that ratios provide an excellent method for determining the temperature and the chemical potential since many theoretical uncertainties cancel out. All indications are that particle abundances are very close to chemical equilibrium. The amount of strangeness present in $S-S$ collisions is very close to the one corresponding to chemical equilibrium. Deviations are at the 30% level.

## 1 Introduction

Considerable efforts have been made to determine the abundance of strange and other particles in ultra-relativistic heavy ion collisions in order to establish the composition of the final state and to infer about the history of the produced system [1, 2, 3]. Due to the complexity of the final state it is natural to attempt a statistical description based on the hadronic gas model. This has been done by several groups recently with considerable success [4, 5, 6, 7, 8, 9, 10, 11].

Since the hadronic gas model uses statistical weight factors there are only very few parameters to describe the freeze-out stage of the produced system. In the Boltzmann approximation the number of particles of type $i$, $N_i^0$, is given by

$$\begin{aligned} N_i^0 &= g_i V \int \frac{d^3p}{(2\pi)^3} e^{-E/T+\mu_i/T} \\ &= \frac{g_i m_i^2 T}{2\pi^2} K_2(m_i/T) e^{\mu_i/T} \end{aligned} \tag{1}$$

where $g_i$ is the degeneracy factor and the index i refers to the particle species under consideration. The corresponding particle density will be denoted by $n_i^0 \equiv N_i^0/V$.

*Talk presented at the "Third International Conference on Physics and Astrophysics of Quark-Gluon Plasma", March 17-21, Jaipur, India.

In the hadron gas model [12] one assumes that all hadronic resonances contribute according to their statistical weight factors. At freeze-out all hadronic resonances decay into the lightest stable particles. All these decays are taken into account by using the known branching ratios.

$$N_{\pi^+} = \sum_i N_i^0 Br(i \to \pi^+) \tag{2}$$

Figure 1 gives an indication of the importance of resonances. In this figure we plot the ratio of thermal pions over the number of all pions (i.e. including those coming from resonance decays). If the temperature is low, resonances are unimportant and the ratio is approximately one. However at a temperature of $T = 200$ MeV this ratio is down to the 10% level. Thus to a good approximation there are no thermal pions if the temperature is above 200 MeV.

For a realistic comparison with experimental data it will be necessary to modify the above expression in several ways: to name but a few, the fireball will not be produced at rest, there will be hard core interactions, there could be a superposition of fireballs, flow effects [2, 3] etc... The expression for the particle number given above will thus have to be modified. We will show below that many of these effects however cancel out when considering hadronic ratios. Thus if $N_i$ is the particle number taking into account various effects, then in many cases one will have the remarkable result

$$\frac{N_i}{N_j} = \frac{N_i^0}{N_j^0} \tag{3}$$

where $N_i$ refers to the particle abundance inclusive of effects due to flow, transverse expansion, excluded volume while $N_i^0$ is the particle abundance calculated in a straightforward manner using Boltzmann statistics without modification, as given above. This is of course not always the case but for several interesting cases which have been widely discussed it is. Therefore hadronic ratios can be used to determine basic thermodynamic quantities like the freeze-out temperature and baryon chemical potential in a way which is free from many distortions (e.g. transverse flow). We will consider in particular the following : 1) excluded volume corrections cancel out if the particle ratios involve an integration over $4\pi$, 2) the effects due to a superposition of fireballs cancel out for ratios provided the abundances are integrated over all values of rapidity, 3) the effects of flow arising from instantaneous freeze-out cancel out if one considers ratios integrated over transverse momenta, 4) for Bjorken longitudinal expansion accompanied by transverse flow, the effects cancel out if one considers particle numbers integrated over all transverse momenta.

Before discussing this in more detail we present quantitatively what is meant by chemical equilibrium.

# 2 Chemical Equilibrium

Hadronic matter has a certain composition of particle species, characterized by "chemical" potentials. In addition to the baryon number chemical potential, $\mu_B$, we have a strangeness chemical potential, $\mu_S$, and also a charge chemical potential $\mu_Q$.. The value of $\mu_B$ is fixed by giving the overall baryon number density $n_B$, and that of $\mu_S$ by fixing the overall strangeness to zero. The value of $\mu_Q$ is fixed by the total baryon number $B$ and the total charge $Q$. If $B/2Q$=1 the system is symmetric in isospin and the charge chemical potential will be zero :





Figure 1: *Ratio of thermal pions to all pions (including pions from resonance decays).*

$\mu_Q = 0$. This is the case for $S - S$ scattering but not for $p - p$ and also not for $Pb - Pb$ where $B/2Q = 0$ and 1.5 respectively. For given values of $T$, $\mu_B$ and $\mu_Q$ one tunes the remaining parameter, $\mu_S$, in such a way as to ensure strangeness neutrality. An example is given in table 1 where we show the relevant Boltzmann factors for different particles in the case of chemical equilibrium. If there is no chemical equilibrium the particle abundances are unrestricted and their normalizations are arbitrary. Dynamical models are then necessary to establish particle abundances [13, 14]. If the system is not in chemical equilibrium than there is no relation between different chemical potentials. In fact the number of chemical potentials necessary to describe the state of the system gives an idea of the degree of equilibration taking place. As perfect chemical equilibrium is of course an extreme case one would like to parametrize deviations from in a simple way. It has been known since a long time that there is a strangeness deficiency in $p - p$ collisions, A qualitative analysis of this was done by Wroblewski [15] many years ago. The deviation from equilibrium can be parameterized by a quantity $\gamma_S$ [16, 17]. Recent estimates [8] give $\gamma_S \approx 0.5$ in $p - p$ collisions. In relativistic heavy ion collisions all estimates lead to a larger value $\gamma_S \approx 0.7 - 1.0$.

We believe that it is very important to establish whether or not chemical equilibrium has been reached.

| | Chemical Equilibrium | No Chemical Equilibrium |
|---|---|---|
| $\pi$ | $\exp\left[-\frac{E_\pi}{T}\right]$ | $\exp\left[-\frac{E_\pi}{T}+\frac{\mu_\pi}{T}\right]$ |
| $N$ | $\exp\left[-\frac{E_N}{T}+\frac{\mu_B}{T}\right]$ | $\exp\left[-\frac{E_N}{T}+\frac{\mu_N}{T}\right]$ |
| $\overline{N}$ | $\exp\left[-\frac{E_N}{T}-\frac{\mu_B}{T}\right]$ | $\exp\left[-\frac{E_N}{T}+\frac{\mu_{\overline{N}}}{T}\right]$ |
| $\Lambda$ | $\exp\left[-\frac{E_\Lambda}{T}+\frac{\mu_B}{T}-\frac{\mu_S}{T}\right]$ | $\exp\left[-\frac{E_\Lambda}{T}+\frac{\mu_\Lambda}{T}\right]$ |
| $\overline{\Lambda}$ | $\exp\left[-\frac{E_\Lambda}{T}-\frac{\mu_B}{T}+\frac{\mu_S}{T}\right]$ | $\exp\left[-\frac{E_\Lambda}{T}+\frac{\mu_{\overline{\Lambda}}}{T}\right]$ |
| $K$ | $\exp\left[-\frac{E_K}{T}+\frac{\mu_S}{T}\right]$ | $\exp\left[-\frac{E_K}{T}+\frac{\mu_K}{T}\right]$ |
| $\overline{K}$ | $\exp\left[-\frac{E_K}{T}-\frac{\mu_S}{T}\right]$ | $\exp\left[-\frac{E_K}{T}+\frac{\mu_{\overline{K}}}{T}\right]$ |

*Table 1 : Boltzmann factors for a hadronic gas with $B/2Q = 1$ in thermal equilibrium.*

# 3 Hadronic Ratios.

## 3.1 Excluded Volume Corrections.

In many cases one includes the hard core repulsive nuclear force in the statistical description by giving hadrons a proper volume. In many cases [18, 19] the particle density is modified in a geometric way leading to the following form

$$\frac{n_i}{n_j} = \frac{\frac{n_i^0}{1+\sum_l n_l^0 V_0}}{\frac{n_j^0}{1+\sum_l n_l^0 V_0}} \tag{4}$$

In this case it is clear that the corrections due to the hard core repulsion cancel out in the ratio of different hadron species and one obtains

$$\frac{n_i}{n_j} = \frac{n_i^0}{n_j^0} \tag{5}$$

This also holds for the excluded volume corrections considered in ref. [19] (provided one uses the Boltzmann approximation) but not for the one considered in ref. [4].

## 3.2 Superposition of Fireballs.

Thermal models based on a single fireball produced in the center-of-mass of the collisions fail to reproduce the observed rapidity distribution. To remedy this one often considers that a distribution of fireballs is being produced. For a symmetric system like $S-S$ or $Pb-Pb$ it is a good start to consider a uniform distribution of fireballs along the rapidity axis but it could be more generally described by a function $\rho(Y_{FB})$ where $Y_{FB}$ is the rapidity of the fireball under consideration. The particle abundance resulting from such a distribution is obtained by integrating over all possible sources. The ratio of two hadronic species is thus given by

$$\frac{n_i}{n_j} = \frac{\int_{-\infty}^{\infty} dy \int_{-Y}^{Y} dY_{FB}\ \rho(Y_{FB}) \frac{dn_i^0}{dy}(y - Y_{FB})}{\int_{-\infty}^{\infty} dy \int_{-Y}^{Y} dY_{FB}\ \rho(Y_{FB}) \frac{dn_j^0}{dy}(y - Y_{FB})} \tag{6}$$

If all the fireballs have the same temperature and the same chemical potential then, after changing the order of integration, one obtains

$$\frac{n_i}{n_j} \quad = \quad \frac{n_i^0}{n_j^0} \frac{\int_{-Y}^{Y}\ dY_{FB}\ \rho(Y_{FB})}{\int_{-Y}^{Y}\ dY_{FB}\ \rho(Y_{FB})} \tag{7}$$

as the integral over fireball distributions is the same in both cases, one obtains again

$$\frac{n_i}{n_j} = \frac{n_i^0}{n_j^0} \tag{8}$$

and the effects cancel out.

## 3.3 Transverse Flow

Recent results from $Pb-Pb$ collisions show that heavy particles like protons and deuterons have a much larger transverse momentum than light particles such as pions and kaons [1, 2, 3]. Within the framework of thermal models this indicates that fireballs have flow in the transverse or at the very least have a random walk type of distribution in the transverse velocity plane. The hydrodynamic description has been extensively discussed in recent papers [20, 21]. The transverse momentum dependence can also be explained by assuming fireballs which have a gaussian distribution in the transverse rapidity. This can be due to either hydrodynamic expansion with flow in the transverse direction or to fireballs being produced with a random walk type of distribution in transverse rapidity [22, 23]. For the discussion about transverse flow it is helpful to have the following two analytic results

$$\int_0^{\infty} dp_T\ p_T m_T^i\ \ K_1\left(\frac{m_T^i}{T}\cosh y_T\right) I_0\left(\frac{p_T}{T}\sinh y_T\right) = m_i^2 T K_2\left(\frac{m^i}{T}\right)\cosh y_T \tag{9}$$

and

$$\int_0^{\infty} dp_T\ p_T p_T\ K_0\left(\frac{m_T^i}{T}\cosh y_T\right) I_1\left(\frac{p_T}{T}\sinh y_T\right) = m_i^2 T K_2\left(\frac{m^i}{T}\right)\sinh y_T \tag{10}$$

### 3.3.1 Instantaneous Freeze-out

If the system undergoes instantaneous freeze-out, the mathematical treatment simplifies considerably since no assumptions have to be made about the freeze-out surface. The velocity profile at freeze-out time can be arbitrary but the system should have the same freeze-out temperature everywhere. The expression for the particle abundance is given by :

$$\frac{n_i}{n_j} = \frac{\int r\ dr \int_0^\infty dp_T m_T^i\ \ K_1\left(\frac{m_T^i}{T}\cosh y_T\right) I_0\left(\frac{p_T}{T}\sinh y_T\right)}{\int r\ dr \int_0^\infty dp_T m_T^j\ \ K_1\left(\frac{m_T^j}{T}\cosh y_T\right) I_0\left(\frac{p_T}{T}\sinh y_T\right)} \tag{11}$$

Using the integrals given above in eqs. (9) and (10), we obtain

$$\frac{n_i}{n_j} = \frac{m_i^2\ \ K_2\left(\frac{m^i}{T}\right)\int r\ dr\ \cosh y_T}{m_j^2\ \ K_2\left(\frac{m^i}{T}\right)\int r\ dr\ \cosh y_T} \tag{12}$$

so that all effects of flow cancel out in the ratio and we are left with the result

$$\frac{n_i}{n_j} = \frac{n_i^0}{n_j^0}. \tag{13}$$

A very similar cancellation also occurs in other models, e.g. the model that was recently considered in ref. [22] based on fireballs having a random walk type of distribution in the transverse direction leads to the same cancellation in the hadronic ratios.

## 3.4 Bjorken scaling + Transverse expansion

If freeze-out is not instantaneous an extra term appears in the expression for the particle spectrum [20, 21]

$$\begin{aligned}\left(\frac{dN_i}{dym_T dm_T}\right)_{y=0} &= \frac{g}{\pi}\int_\sigma r\ dr\ \tau_F(r) \\ &\quad \left\{m_T^i I_0\left(\frac{p_T\sinh y_T}{T}\right) K_1\left(\frac{m_T^i\cosh y_T}{T}\right)\right. \\ - &\quad \left.\left(\frac{\partial\tau_F}{\partial r}\right) p_T I_1\left(\frac{p_T\sinh y_T}{T}\right) K_0\left(\frac{m_T^i\cosh y_T}{T}\right)\right\}\end{aligned} \tag{14}$$

where the partial derivative $\partial\tau_F/\partial r$ has to be taken on the freeze-out surface, denoted by $\sigma$ in the equation above. After integration over $m_T$ (and only after this integration) one obtains

$$\begin{aligned}\left(\frac{dN_i}{dy}\right)_{y=0} = &\quad \frac{g}{\pi}\int_\sigma r\ dr\ \tau_F(r) \\ &\quad \left\{\cosh(y_T) - \left(\frac{\partial\tau_F}{\partial r}\right)\sinh(y_T)\right\} m_i^2 T K_2\left(\frac{m_i}{T}\right)\end{aligned} \tag{15}$$





Figure 2: *Particle ratios using data from the AGS at BNL.*

leading to the result

$$\frac{dN_i/dy}{dN_j/dy} = \frac{N_i^0}{N_j^0} \tag{16}$$

This result is remarkable : in a model where the transverse spectra are severely distorted by transverse flow and the rapidity distribution has a plateau, the particle ratios are still given by the uncorrected Boltzmann distribution as if particles originated from a fireball at rest. Effects of hydrodynamic flow cancel out in the ratio provided the particle abundance has been integrated over transverse momentum.

# 4 Results

The above discussion has given us some confidence that the particle ratios can be used to determine the temperature and the chemical potential in a given reaction. Comparison of hadronic abundances with thermal models has been done recently for $p-p$ collisions [8], for $Si-Au$ [4, 7] and $Au-Au$ [5] collisions at BNL, and also for the preliminary results of the results from $Pb-Pb$ at CERN. Each particle ratio leads to a band in the $T, \mu_B$ plane, (including the error bar) this gives rise to a region where the hadronic gas reproduces the measured numbers. If all the measured ratios have a common overlap region then one can conclude that the experimental data are consistent with chemical equilibrium. The data from Brookhaven lead to $T \approx 100-120$ Mev and a chemical potential which is surprisingly well determined $\mu_B$=540





Figure 3: *Strange particle ratios using data from the WA85 collaboration.*

MeV. The CERN data give $T \approx 180$ MeV and a chemical potential which is substantially lower than the one from BNL $\mu_B \approx$ 200-300 MeV. A discussion of different break-up mechanisms has been presented in [24, 25, 26].

# 5 Conclusions

One of the important parameters in the description of relativistic heavy ion collisions is the temperature. We have argued that the best way to determine directly the temperature is to consider ratios of hadronic yields that have been integrated over transverse momentum. In such a ratio many effects (e.g. flow) cancel out. A surprisingly large number of hadronic abundances can be explained in this way with a temperature which is around 180 MeV for data obtained from $S - S$ collisions at CERN and around 110 MeV for data from the AGS. One conclusion forces itself upon us : there is nothing strange about strangeness.

## Acknowledgment

We gratefully acknowledge useful and stimulating discussions with Helmut Satz, Krzysztof Redlich and Dinesh Srivastava. The financial support of the University of Cape Town (URC) and of the Foundation for Research Development (FRD) are gratefully acknowledged.

# References


[1] *"Quark Matter '96"*, Proc. of the XII'th Int. Conf. on Ultra-Relativistic Nucleus-Nucleus Collisions, Heidelberg, 1996, P. Braun-Munzinger, H.J. Specht, R. Stock and H. Stöcker (editors), Nucl. Phys. **A610** (1996).

[2] L.G. Bearden et al. (NA44 collaboration), Phys. Rev. Letters **78**, 2080 (1997).

[3] P. Jacobs at al. (NA49 collaboration) contribution to this conference.

[4] P. Braun-Munzinger, J. Stachel, J. P. Wessels, and N. Xu, Phys. Lett. **B344**, 43 (1995); B **365**, 1 (1996).

[5] P. Braun-Munzinger and J. Stachel, Nucl. Phys. **A606**, 320 (1996).

[6] F. Becattini, Z. f. Physik **C69**, 485 (1996).

[7] J. Cleymans, D. Elliott, H. Satz and R.L. Thews, Z. f. Physik **C74**, 319 (1997).

[8] F. Becattini and U. Heinz, Firenze preprint DFF 268/02/1997, Z. f. Physik **C** to be published.

[9] J. Letessier, A. Tounsi, U. Heinz, J. Sollfrank and J. Rafelski, Phys. Rev. **D51**, 3408 (1995).

[10] J. Sollfrank et al., Z. f. Physik **C61**, 659 (1994).

[11] M.N. Asprouli and A.D. Panagiotou, Phys. Rev. **C51**, 1444 (1995).

[12] See e.g. J. Cleymans and H. Satz, Z. f. Physik **C57**, 135 (1993).

[13] H.W. Barz, B.L. Friman, J. Knoll and H. Schulz, Nucl. Phys. **A484**, 661 (1988).

[14] C. Spieles, H. Stöcker and C. Greiner, report nucl-th/9704008.

[15] A. Wroblewski, Acta Phys. Pol. **B16**, 379 (1985).

[16] J. Rafelski, Nucl. Phys. **A544**, 279c (1992).

[17] C. Slotta, J. Sollfrank and U. Heinz, in *"Strangeness in Hadronic Matter."*, Ed. J. Rafelski, AIP Conference Proceedings 340, 1995.

[18] J. Cleymans, K. Redlich, H. Satz and E. Suhonen, Z. f. Physik **C33**, 151 (1986).

[19] D.K. Rischke, M. Gorenstein, H. Stöcker and W. Greiner, Z. f. Physik, **C51**, 485 (1991).

[20] J. Cleymans, K. Redlich, and D. K. Srivastava, Phys. Rev. **C55**, 1431 (1997).

[21] J. Sollfrank et al. Phys. Rev. C55 1431 (1997).

[22] A. Leonidov, M.Nardi and H. Satz, Z. f. Physik **C74**, 535 (1997).

[23] S. Esumi, U. Heinz and N. Xu, preprint March 1997.

[24] K. Redlich, J. Cleymans, H. Satz and E. Suhonen, Nucl. Phys. **A566**, 391c (1994).

[25] U. Heinz, Nucl. Phys. **A566** 205c (1993).

[26] J. Cleymans, K. Redlich, H. Satz and E. Suhonen, Z. f. Physik **C58**, 347 (1993).